

# <sup>10</sup>Be ages of late Pleistocene deglaciation and Neoglaciation in the north-central Brooks Range, Arctic Alaska



MICHAEL E. BADDING,<sup>1</sup> JASON P. BRINER<sup>1\*</sup> and DARRELL S. KAUFMAN<sup>2</sup>

<sup>1</sup>Department of Geology, State University of New York at Buffalo, Buffalo, NY 14260, USA

<sup>2</sup>School of Earth Sciences & Environmental Sustainability, Northern Arizona University, Flagstaff, AZ 86011, USA

Received 28 June 2012; Revised 18 September 2012; Accepted 20 September 2012

**ABSTRACT:** We present a chronology of late Pleistocene deglaciation and Neoglaciation for two valleys in the north-central Brooks Range, Alaska, using cosmogenic <sup>10</sup>Be exposure dating. The two valleys show evidence of ice retreat from the northern range front before ~16–15 ka, and into individual cirques by ~14 ka. There is no evidence for a standstill or re-advance during the Lateglacial period, indicating that a glacier advance during the Younger Dryas, if any, was less extensive than during the Neoglaciation. The maximum glacier expansion during the Neoglacial is delimited by moraines in two cirques separated by about 200 km and dated to  $4.6 \pm 0.5$  and  $2.7 \pm 0.2$  cal ka BP. Both moraine ages agree with previously published lichen-inferred ages, and confirm that glaciers in the Brooks Range experienced multiple advances of similar magnitude throughout the late Holocene. The similar extent of glaciers during the middle Holocene and the Little Ice Age may imply that the effect of decreasing summer insolation was surpassed by increasing aridity to limit glacier growth as Neoglaciation progressed. Copyright © 2012 John Wiley & Sons, Ltd.

**KEYWORDS:** Alaska; Brooks Range; Neoglaciation; <sup>10</sup>Be dating.

## Introduction

Arctic environments are affected by cryosphere–albedo feedback mechanisms that result in climate amplification (Overpeck *et al.*, 1997; Holland and Bitz, 2003; Serreze and Barry, 2011), and the Arctic is expected to undergo significant change during the 21st century (Miller *et al.*, 2010; Sharp *et al.*, 2011; Jacob *et al.*, 2012). Glaciers are one component of the Arctic environment, and have experienced pronounced retreat due to recent climate warming (Meier *et al.*, 2003; Gardner *et al.*, 2011; Fisher *et al.*, 2012). Reconstructions of past glacier fluctuations can provide insights into Arctic climate change. In particular, alpine glaciers respond sensitively to climate change, and when well-dated records of former glacier change are available from around the globe, they can be used to reconstruct spatial patterns of climate change. However, there are a limited number of well-dated alpine glacier chronologies in the Arctic.

The Brooks Range, Arctic Alaska (Fig. 1), affords an opportunity to investigate the timing of alpine glacier retreat because unlike most land area in the Arctic, the Brooks Range was glaciated by alpine glaciers rather than covered by ice sheets. Determining the timing of deglaciation in the Brooks Range provides a means to investigate Arctic climate patterns during the last termination, and to compare these with glacier records from elsewhere in the northern hemisphere. For example, it is unknown whether glaciers in the Brooks Range responded to the abrupt climate changes that define the Lateglacial period in the North Atlantic region (Alley *et al.*, 1993; Denton *et al.*, 2010; Harrison *et al.*, 2010). Although glacier advance during the Younger Dryas period (YD; 12.9–11.7 ka) may be prevalent in some areas around the North Atlantic region (Ingólfsson *et al.*, 1997; Lohne *et al.*, 2012), evidence elsewhere is less definitive (e.g. Gosse *et al.*, 1995; Mangerud and Landvik, 2007; Kelly *et al.*, 2008).

The Brooks Range contains ~1000 modern glaciers, which have been retreating significantly in recent decades (Molina,


2007). It is unknown when glaciers in the Brooks Range expanded following the Holocene warm period (Kaufman *et al.*, 2004), and when they achieved their maximum extent during the Holocene. Previous studies suggest that numerous mid-Holocene moraines are preserved outboard of moraines deposited during the Little Ice Age (LIA; ~1200–1900 AD; Calkin and Ellis, 1980; Ellis and Calkin, 1984; Solomina and Calkin, 2003; Sikorski *et al.*, 2009). Pre-LIA late Holocene moraines are absent in front of many glaciers throughout the northern hemisphere, probably because most glaciers seem to have been driven by the decrease in northern latitude summer insolation throughout the Holocene and they culminated in their greatest extent during the LIA (e.g. Barclay *et al.*, 2009; Briner *et al.*, 2009). Although the mechanism behind Arctic summer cooling throughout the Holocene is well understood on millennial timescales, dating the multiple glacier fluctuations throughout the late Holocene provides additional information about centennial-scale climate changes in the Arctic.

The goals of this study are to: (i) develop a chronology of final deglaciation during the last termination; and (ii) produce numerical ages on outermost late Holocene moraines. We present 17 cosmogenic <sup>10</sup>Be exposure ages (hereafter termed <sup>10</sup>Be ages) on deglacial and Holocene glacial landforms in the north-central Brooks Range. We report <sup>10</sup>Be ages from the headwaters of two valleys to determine when the upper catchments became deglaciated. In addition, we <sup>10</sup>Be-dated two pre-LIA moraine crests with previously published lichenometric ages to determine when Brooks Range glaciers attained their maximum Neoglacial configuration.

## Setting and glacial history

The Brooks Range is the northernmost mountain range in Alaska, trending east–west and separating the North Slope to the north from the Yukon River basin to the south (Fig. 1). The range stretches ~1000 km from the Chukchi Sea on the west coast to the Alaska–Yukon border in the east, and lies entirely north of the Arctic Circle. The Brooks Range reaches over 2700 m above sea level (a.s.l.), with ~1000 glaciers (mostly

\*Correspondence: J. P. BRINER, as above.  
E-mail: jbriner@buffalo.edu



**Figure 1.** Digital elevation model of north-central Brooks Range. Inset: location of Brooks Range within eastern Beringia. Locations mentioned in text are shown; RMV, Roche Moutonnée valley. Ice limits from Hamilton (1986). This figure is available in colour online at [wileyonlinelibrary.com](http://wileyonlinelibrary.com).

cirque glaciers) occupying the highest peaks (Molina, 2007). Nearly all of the modern glaciers are sheltered behind north-facing cirques (Ellis, 1982). Bedrock in the north-central Brooks Range is dominated by highly deformed Devonian sedimentary and meta-sedimentary strata (Brosgé *et al.*, 1979). Most relevant for  $^{10}\text{Be}$  dating are the resistant, quartz-rich members of the Kanayut Conglomerate (Upper Devonian). Although Pleistocene drift limits are well mapped in the Brooks Range, and the chronology of several Itkillik II maxima phases (late Wisconsin glaciation) is generally known (Hamilton, 2003; Balascio *et al.*, 2005; Briner and Kaufman, 2008), the timing of deglaciation in the Brooks Range is less well known (Hamilton, 1986).

We collected samples for  $^{10}\text{Be}$  dating to constrain the timing of deglaciation of the headwaters in the Kurupa River and Atigun River valleys. We chose these study locations because both valleys are accessible, contain the target lithology for  $^{10}\text{Be}$  dating, and were previously mapped and dated with lichenometry (Ellis and Calkin, 1984; Lamb, 1984). The Kurupa River valley is situated in the west-central Brooks Range (Fig. 1). Here, Itkillik II terminal moraines lie  $\sim 10$  km north of the range front and  $\sim 25$  km downvalley of cirque headwalls (Hamilton, 1982). Although the glacial chronology has not been previously studied in the Kurupa River valley, radiocarbon ages ranging from 12.6 to 15.4 cal ka BP from the Okokmilaga River valley,  $\sim 60$  km east (Fig. 1), provide minimum limiting ages on timing of deglaciation from Itkillik II ice limits (Hamilton, 1982).

Approximately 200 km to the east of Kurupa River valley is the Atigun River valley (Fig. 1), which served as a major trough for ice drainage during the last glaciation (Hamilton, 1986). We investigated the deglaciation of Roche Moutonnée valley (informal name, Fig. 1), which is a tributary to the Atigun River valley  $\sim 15$  km south of the northern range front.


Terminal moraines deposited by glaciers sourced in the Atigun River valley during the early phase of the Itkillik II glaciation are located  $\sim 30$  km from the cirque headwalls of Roche Moutonnée valley (Fig. 1). During the final phase of the Itkillik II glaciation in the Atigun River valley, ice spilled eastward through Atigun Gorge and into the Sagavanirktok

River valley (Fig. 1), forming a piedmont lobe and partially damming the Sagavanirktok River (Hamilton, 2003). Radiocarbon ages dating an aggradation event associated with an ice advance into the Sagavanirktok River valley are  $15.2 \pm 1.4$  and  $15.5 \pm 1.2$  cal ka BP (Fig. 1); an additional radiocarbon age of  $13.8 \pm 0.7$  cal ka BP is interpreted as a minimum limiting age on deglaciation (Fig. 1; Hamilton, 2003).

Our Holocene glacial chronology arises from two cirques located  $\sim 200$  km apart, both of which contain pre-LIA moraines based on previous mapping (Ellis and Calkin, 1984; Lamb, 1984). In the Kurupa River valley, we collected samples from boulder surfaces from the outermost pre-LIA moraine fronting Fireweed West glacier [all glacier names are informal from Ellis and Calkin (1984) and Lamb (1984)]; previous mapping and lichenometric data suggested that a suite of moraines stabilized at  $750 \pm 150$ ,  $1650 \pm 330$  and  $3700 \pm 740$  cal a BP (Lamb, 1984). We also dated moraine boulders from the Triple East glacier, which hosts one of the oldest lichenometry-inferred Holocene moraines in the north-central Brooks Range (Ellis and Calkin, 1984; Fig. 1). Ellis and Calkin (1984) assigned lichen ages of  $400 \pm 80$ ,  $1900 \pm 380$ ,  $3200 \pm 640$ ,  $3700 \pm 740$  and  $5000 \pm 1000$  cal a BP to the moraines fronting Triple East glacier.

## Methods

Sites were selected for  $^{10}\text{Be}$  dating by considering potential snow and debris cover and post-glacial erosion. We sampled bedrock and boulder surfaces marked by glacial polish and striations, indicating negligible post-glacial erosion. Samples from bedrock were collected from at least 0.5 m above surrounding landscape and from elevated sites within the valley bottom, increasing the likelihood of the site to be windswept of snow and free of previous cover by glacial sediment (Fig. 2). We sampled sites along the valley center where the duration of glacier occupation and efficiency of subglacial erosion are most likely to have reduced the possibility of isotopic inheritance. In some cases we sampled



**Figure 2.** Examples of boulder and bedrock sample sites. (A) Bedrock knob (sample 11RMV-08) in the middle of the valley and ~40 m above the valley bottom. (B) Perched erratic (11RMV-07) on glacially eroded bedrock (11RMV-08). (C) Typical Holocene moraine boulder (10FWW-02). (D) Bedrock sample site (10KRV-09) with late Holocene moraine in the background. This figure is available in colour online at [wileyonlinelibrary.com](http://wileyonlinelibrary.com).

erratic boulders that rested directly on exposed bedrock (Fig. 2). All moraine boulders that we sampled had tops ~1 m above the surrounding landscape and had no evidence of post-depositional movement (Fig. 2). The Fireweed West glacier moraines are probably ice cored, and thus some downslope movement is likely to have occurred. However, we avoided portions of the moraine with boulders that exhibited obvious signs of rotation, aided with lichenometry. Thus, despite some post-depositional down-valley motion of the moraines, many boulders on the moraine crests are probably in the same orientation today as they were when they were deposited originally. All samples were from horizontal to near-horizontal surfaces, and we avoided boulder edges and corners. Geographic coordinates and elevation for each sample were determined with a handheld GPS device, and checked on a 1:63 360-scale USGS topographic map. Topographic shielding was measured in the field at each sample site with a clinometer and compass.

Samples were collected with a hammer and chisel from ice-sculpted meta-sandstone and conglomerate bedrock surfaces ( $n=8$ ), erratics perched on bedrock ( $n=2$ ) and moraine boulders ( $n=7$ ). All samples were processed at the University at Buffalo Cosmogenic Isotope Laboratory. Samples were crushed and sieved to isolate the 250- to 425- $\mu\text{m}$  size fraction. Samples were pretreated in dilute HCl and HNO<sub>3</sub>-HF acid baths. Quartz was isolated with a series of heated HNO<sub>3</sub>-HF acid baths, and in some cases large grains with visible inclusions were removed by hand. <sup>9</sup>Be carrier (~0.37–0.80 mg; 405 p.p.m.) was added to each sample prior to dissolution in concentrated HF. Beryllium was extracted using ion-exchange chromatography, selective precipitation with NH<sub>4</sub>OH and final oxidation to BeO.

<sup>10</sup>Be/<sup>9</sup>Be accelerator mass spectrometry measurements were completed at the Lawrence Livermore National Laboratory Center for Mass Spectrometry and normalized to standard 07KNSTD3110 (Nishiizumi *et al.*, 2007). Ratios for process blanks ( $n=2$ ) averaged  $4.39 \pm 2.06 \times 10^{-15}$ . <sup>10</sup>Be ages were calculated assuming no snow shielding and no erosion using the CRONUS-Earth online exposure age calculator (<http://hess.ess.washington.edu/math>; Version 2.2; Balco *et al.*, 2008).

We used the north-eastern North America <sup>10</sup>Be production rate of  $3.93 \pm 0.19$  atoms  $\text{g}^{-1} \text{a}^{-1}$  (Balco *et al.*, 2009) and the constant-production scaling scheme of Lal (1991) and Stone (2000). There are no <sup>10</sup>Be production-rate calibration sites in Alaska; thus, we choose the north-eastern North America <sup>10</sup>Be production rate because recent <sup>10</sup>Be production rates from elsewhere in the Arctic yield a similar value (Fenton *et al.*, 2011; Briner *et al.*, 2012). The north-eastern North American <sup>10</sup>Be production rate yields ages ~12% older than the globally averaged rate from Balco *et al.* (2008) when using the 'St' scaling scheme. The use of alternative scaling schemes results in ages that are up to ~4.9% older or ~1.5% younger than the ages that we present.

## Results

### Late Pleistocene deglaciation


We <sup>10</sup>Be-dated five samples from four sites in the eastern portion of upper Kurupa River valley (Fig. 1). Four <sup>10</sup>Be ages from valley-bottom bedrock samples and one <sup>10</sup>Be age from an erratic boulder range from  $15.9 \pm 0.5$  to  $13.8 \pm 0.3$  ka (Table 1, Fig. 3), and span 1.5–6.5 km from the cirque headwall. The ages increase with distance from cirque headwalls. Samples from the erratic boulder and bedrock upon which it lies yield statistically identical <sup>10</sup>Be ages of  $13.9 \pm 0.3$  ka (erratic) and  $13.8 \pm 0.3$  ka (bedrock). Five <sup>10</sup>Be ages from four sites within Roche Moutonnée valley range in age from  $50.4 \pm 1.2$  to  $14.7 \pm 0.5$  ka (Table 1, Fig. 4) spanning 3–10 km from the cirque headwall. Although the ages in Roche Moutonnée valley are considerably older than in Upper Kurupa valley, they too decrease in age toward the headwalls. Samples from an erratic boulder and bedrock upon which it is perched yield <sup>10</sup>Be ages of  $14.7 \pm 0.5$  and  $17.1 \pm 0.3$  ka, respectively.

### Mid Holocene moraines

In the upper Kurupa River valley, we adopted the Holocene moraine mapping from Lamb (1984), who showed an end-moraine complex fronting Fireweed West glacier consisting of


**Table 1.** <sup>10</sup>Be ages, sample data and location information.

Sample	Sample type	Latitude (°N) (DD)	Longitude (°W) (DD)	Elevation (m a.s.l.)	Sample height (m)	Thickness (cm)	Shielding correction	Quartz (g)	Be carrier added (g)	<sup>10</sup> Be/ <sup>9</sup> Be	<sup>10</sup> Be uncertainty (atoms g <sup>-1</sup> )	<sup>10</sup> Be age (ka)
<b>Kurupa River Valley</b>												
10FWW-01	Moraine boulder	68.19317	154.51979	1499	1.5	1.5	0.979	30.0549	0.3687	1.421E-13	4.150E-15	2.8 ± 0.1
10FWW-02	Moraine boulder	68.19309	154.51994	1508	1.0	1.0	0.979	35.0691	0.3704	1.589E-13	2.373E-15	2.6 ± 0.1
10FWW-03	Moraine boulder	68.19267	154.51959	1509	1.0	1.0	0.979	35.5775	0.3718	1.335E-13	3.392E-15	2.2 ± 0.1
10KRV-03	Bedrock	68.22772	154.50264	1267	0.5	2.0	0.993	29.3285	0.8008	2.904E-13	8.403E-15	15.9 ± 0.4
10KRV-05	Bedrock	68.21104	154.49136	1296	2.0	2.0	0.983	35.0942	0.8029	3.368E-13	6.296E-15	15.2 ± 0.3
10KRV-07	Erratic Boulder	68.20375	154.48836	1318	1.5	2.0	0.972	35.2659	0.3711	6.702E-13	1.241E-14	13.8 ± 0.3
10KRV-08	Bedrock	68.20383	154.48865	1314	1.5	3.0	0.972	35.0602	0.3722	6.596E-13	1.218E-14	13.9 ± 0.3
10KRV-09	Bedrock	68.19965	154.52574	1476	1.0	3.0	0.981	35.0313	0.3724	7.570E-13	1.408E-14	13.8 ± 0.3
<b>Atigun River Valley region</b>												
11TE-01	Moraine boulder	68.33395	149.76705	1846	1.0	2.0	0.993	40.0375	0.8035	1.616E-13	4.755E-15	4.1 ± 0.1
11TE-02	Moraine boulder	68.33423	149.76735	1849	0.75	2.5	0.993	30.0701	0.7962	1.545E-13	6.141E-15	5.1 ± 0.2
11TE-03	Moraine boulder	68.33510	149.76785	1833	1.5	2.5	0.993	30.0480	0.8031	1.361E-13	2.526E-15	4.6 ± 0.1
11TE-04	Moraine boulder	68.33532	149.76832	1826	1.7	1.5	0.993	35.1430	0.7982	9.146E-14	2.618E-15	2.6 ± 0.1
11RMV-02	Bedrock	68.30927	149.14107	1246	1.5	4.0	0.983	35.0755	0.8014	3.238E-13	5.990E-15	15.6 ± 0.3
11RMV-07	Erratic Boulder	68.33232	149.15842	1242	1.0	1.0	0.990	32.9613	0.7965	2.975E-13	9.993E-15	14.7 ± 0.5
11RMV-08	Bedrock	68.33235	149.15790	1235	1.0	1.5	0.990	35.1294	0.7958	3.644E-13	6.764E-15	17.1 ± 0.3
11RMV-13	Bedrock	68.36522	149.25767	1019	2.0	1.5	0.995	32.2053	0.8042	3.630E-13	6.732E-15	22.6 ± 0.4
11RMV-15	Bedrock	68.38003	149.31113	876	0.5	1.0	0.996	35.0198	0.8037	7.759E-13	1.759E-14	50.4 ± 1.2




**Figure 3.** Aerial photograph of the headwaters of the Kurupa River valley. The Itkillik II ice limit was ~25 km north downvalley. Numbers are <sup>10</sup>Be ages reported in cal ka BP. LIA moraines are outlined by white dashed lines. USGS aerial photograph acquired in 1982. LG, Lupine glacier; CG, Cotton Grass glacier; FWWG, Fireweed West glacier; UPK, Upper Kurupa Lake.

multiple ridges, the outermost of which has lichen diameters consistent with pre-LIA deposition (Supporting Information, Figs S1 and S2). Our <sup>10</sup>Be analyses on three moraine boulders from the outermost moraine fronting this glacier yield ages of 2.2 ± 0.1, 2.6 ± 0.1 and 2.8 ± 0.1 ka (Table 1, Fig. 5).



**Figure 4.** Aerial photograph of the Atigun River valley and headwaters of the Roche Moutonnée valley. The Itkillik II ice limit was 15 km downvalley (north). <sup>10</sup>Be ages are reported in cal ka BP. USGS aerial photograph acquired in 1978.



**Figure 5.** Vertical views of Fireweed West and Triple East glaciers showing Neoglacial moraine crests and locations of samples for  $^{10}\text{Be}$  ages (reported in cal ka BP). USGS aerial photographs acquired in 1982 (Fireweed West), and Triple East Glacier is shown in a Google Earth image acquired on 19 September 2010. This figure is available in colour online at [wileyonlinelibrary.com](http://wileyonlinelibrary.com).

dimensions range from  $2 \times 1.5 \times 1.5$  m for the smallest boulder to  $2 \times 2 \times 1.5$  m for the largest boulder.

At Triple East Glacier, we adopted the mapping from Ellis and Calkin (1984), who showed five moraine crests on the north-eastern portion of the latero-end moraine complex (supporting Fig. S3). We sampled four boulders from the outermost of these moraine crests, which yield ages of  $2.6 \pm 0.1$ ,  $4.1 \pm 0.1$ ,  $4.6 \pm 0.1$  and  $5.1 \pm 0.2$  ka (Table 1, Fig. 5). Boulder dimensions range from  $1 \times 1 \times 1$  m for the smallest boulder to  $2.5 \times 2 \times 1.7$  m for the largest boulder.

## Interpretation and discussion

### Deglaciation

In the upper Kurupa River valley, the  $^{10}\text{Be}$  ages suggest that deglaciation of the range front occurred prior to  $\sim 16$  ka. Glaciers retreated into the upper catchment by 15.9 ka, and ice retreated to within Neoglacial limits, or perhaps disappeared entirely, around 13.8 ka. The youngest age on bedrock of  $13.8 \pm 0.3$  ka, from just beyond the outermost Holocene moraine, implies that the cirques were ice-free or mostly ice-free at this time (Fig. 2). We believe these ages are not influenced by inheritance because they decrease upvalley and the paired erratic and bedrock samples yield identical ages. The timing of deglaciation of the Upper Kurupa River valley is consistent with the minimum radiocarbon age from Okokmilaga valley of  $15.3 \pm 1.8$  cal ka BP (Hamilton, 1986). Glacier retreat appears to have slowed between 15.2 and 13.8 cal ka BP


because of the close proximity of these ages, although no geomorphic evidence (e.g. moraine) suggests a standstill or re-advance in this vicinity.

In the Roche Moutonnée valley, several of the valley-bottom bedrock ages are apparently too old to represent the timing of deglaciation because they are inconsistent with previously published radiocarbon ages (Hamilton, 1986, 2003). We attribute the older-than-expected  $^{10}\text{Be}$  ages in Roche Moutonnée valley to inheritance. We interpret the ages of samples 11RMV-8 ( $17.1 \pm 0.3$  ka), 11RMV-13 ( $22.6 \pm 0.4$  ka) and 11RMV-15 ( $50.4 \pm 1.2$  ka) to be influenced by inheritance (Table 1); although we have more confidence in the ages of 11RMV-2 ( $15.6 \pm 0.3$  ka) and 11RMV-7 ( $14.7 \pm 0.5$  ka) because they are similar to independent chronologies in the Brooks Range of when deglaciation occurred, we cannot rule out completely that they lack inheritance. Minimum limiting radiocarbon ages from Hamilton (1986) suggest ice was perhaps 10 km downvalley in Atigun River valley at  $\sim 15.5$  ka cal BP. If the age of  $14.7 \pm 0.5$  ka from an erratic is not influenced by inheritance, it implies rapid deglaciation at  $\sim 15$  ka.

Inherited cosmogenic isotopes could be the result of partially cold-based glaciers with low rates of subglacial erosion that led to incomplete resetting of the bedrock surface. In addition, the Kanayut Conglomerate, including the beds that we sampled in the Roche Moutonnée valley (Fig. 2), is known to be particularly erosion resistant (Brosgé *et al.*, 1979), whereas the siliclastic and arkosic sandstones that we sampled from the Kurupa River valley are more easily eroded. Differences in erodability may explain the differences in inheritance between the two valleys.

### Regional comparison

Previously published glacier chronologies in the Brooks Range indicate an early maximum extent and early retreat from late Pleistocene positions compared with elsewhere in Alaska (Briner and Kaufman, 2008). However, radiocarbon ages of  $\sim 15$  cal ka BP are only minimum ages on late Pleistocene moraine abandonment at the range front (Hamilton, 1982, 1986). Previous  $^{10}\text{Be}$  dating in the north-eastern Brooks Range



**Figure 6.** Glacier time–distance diagram for the two study valleys. The  $^{10}\text{Be}$  ages from the Kurupa River valley suggest that glaciers had retreated into the cirques prior to the Younger Dryas (YD). Glacier expansion was underway during the middle Holocene, based on data from Triple East (TE) glacier. Other late Holocene glacier fluctuations are based on moraine mapping and lichenometry by Ellis and Calkin (1984) and Lamb (1984). The Greenland isotope record from Stuiver and Grootes (2000) and summer insolation from Berger and Loutre (1991) are shown for comparison. This figure is available in colour online at [wileyonlinelibrary.com](http://wileyonlinelibrary.com).


revealed glaciers there reached their maximum at ~30–27 ka, and had retreated to within ~12 km of their cirque headwalls prior to ~21 ka (ages adjusted according to the production rate used here) (Balascio *et al.*, 2005; Briner *et al.*, 2005). Our <sup>10</sup>Be ages from the Kurupa River valley indicate that ice had retreated to within ~5 km of their cirque headwalls prior to ~16 ka and had retreated into their cirques by ~14 ka (Fig. 6). In southern Alaska, some glaciers were still expanded far downvalley ~16 ka (Briner and Kaufman, 2008). Evidence that glaciers were well downvalley at ~17–16 ka in many south and central Alaska valleys, while at the same time glaciers were within a few kilometers of their cirques in the Brooks Range, reflects the spatial heterogeneity in Alaskan climate.

Our <sup>10</sup>Be ages provide the opportunity to investigate potential Lateglacial advances as observed elsewhere in north hemisphere, such as during the YD. The <sup>10</sup>Be age of 13.8 ± 0.3 ka from the upper Kurupa River valley immediately outboard of the Neoglacial moraines of Cottongrass Glacier (Figs 2 and 3) suggests that any advance during the YD was less extensive than advances during Neoglaciation. Previous studies in the Brooks Range failed to identify glacier advances during the YD (e.g. Hamilton, 2003; Balascio *et al.*, 2005). The lack of evidence for a glacier advance during the YD in the Brooks Range is compatible with findings from Kokorowski *et al.* (2008), who showed similar-to-present or even higher-than-present temperatures across central and northern Alaska during the YD. Cooling during the YD is more strongly expressed in southern Alaska, but was heterogeneous (Kaufman *et al.*, 2010). Elsewhere in the Arctic, including Baffin Island and Svalbard, glaciers were smaller than during the LIA (Mangerud and Landvik, 2007; Briner *et al.*, 2009). On the other hand, isotope records from permafrost on the North Slope indicate cooling during the YD (Meyer *et al.*, 2010), but this is interpreted to reflect lower winter temperatures that might not have led to significant glacier growth. On the Arctic Foothills, lake levels fell and streams incised their floodplains during the YD (Mann *et al.*, 2002), further indication of lowered effective moisture at that time.

### Neoglaciation

At both Fireweed West and Triple East glaciers, <sup>10</sup>Be ages on the outermost moraine include one age noticeably younger than the others (Table 1). Averaging the closely spaced ages and excluding the youngest age from each dataset yields ages of 2.7 ± 0.2 ka (*n* = 2) at Fireweed West glacier and 4.6 ± 0.5 ka (*n* = 3) at Triple East glacier (Fig. 6). Alternatively, we could use the oldest age as a best estimate of the moraine emplacement age, assuming that there was no inheritance and that moraine degradation was the main cause for the age spread (Heyman *et al.*, 2010; Applegate *et al.*, 2012). However, given that these moraines were deposited close to cirque headwalls and contain abundant angular blocks that were probably transported supraglacially, we hesitate to rule out inheritance. Alternatively, the moraines are composite features deposited during multiple periods when glaciers expanded onto their former end moraines, and each <sup>10</sup>Be age represents the timing of deposition of that particular boulder on the moraine crest. Although the tightly nested ridges of the end moraine complex suggest that, as a whole, it is a composite feature, we hypothesize that each ridge crest was built during a single maximum. Regardless, despite the difficulty in interpreting the <sup>10</sup>Be ages, we favor the interpretation that the average of the oldest ages is the best approximation for the age of moraine stabilization.

We compare our <sup>10</sup>Be chronology with the previously published moraine ages determined from lichenometry (Fig. 7).



**Figure 7.** Growth curve for *Rizocarpon geographicum* (*sensu lato*) and <sup>10</sup>Be ages associated with lichenometrically dated moraines (modified from Calkin and Ellis, 1980; Calkin and Solomina, 2003). Gray lines bracketing the growth curve are ±20% uncertainty. \*Assigned moraine age is the average of two ages, excluding the youngest age. Error bars show 2-sigma range.

Lichenometry has an estimated ~20% age uncertainty, and calibration points are limited to the last millennium (Calkin and Ellis, 1980; Solomina and Calkin, 2003). Using the lichen growth curve from Ellis and Calkin (1984), which was converted to calendar years by Solomina and Calkin (2003), our <sup>10</sup>Be ages on Holocene moraines overlap with the 20% uncertainty of the lichenometric age assignments. For example, Lamb (1984) reported lichen diameters that yield an age of 3.7 ± 0.7 ka cal BP on the Fireweed West moraine, and our average <sup>10</sup>Be age is 2.7 ± 0.2 ka. Our average <sup>10</sup>Be age of the Triple East moraine of 4.6 ± 0.5 ka also agrees with the previously estimated age of 5.0 ± 1.0 ka cal BP (Ellis and Calkin, 1984). Based on these <sup>10</sup>Be ages, it seems that the lichen growth curve developed by Ellis and Calkin (1984) can be reliably projected beyond the control points using a linear function as they had suggested, at least to the middle Holocene. The exponential form of the growth curve that was considered by Solomina and Calkin (2003) yields ages that are much too old relative to the <sup>10</sup>Be ages. The second-order polynomial form that was suggested by Sikorski *et al.* (2009) yields ages that are intermediate between the linear and exponential fits.

These <sup>10</sup>Be ages demonstrate that glaciers in the Brooks Range experienced advances in the middle Holocene that were as extensive as during the LIA. In part, the pre-LIA moraine preservation may be due to the inability of smaller subsequent glaciers to overrun previously emplaced moraines (Porter, 2007). Furthermore, those moraine complexes that are ice cored might have post-depositional, net downslope movement, and thus escape subsequent disturbance; however, many moraine sequences throughout the Brooks Range also lack ice cores. Regardless, glaciers during the middle Holocene were at least similar in size to those during the LIA. This is at odds with insolation as the main driver of glacier growth, which seems to explain the fact that glaciers were most extensive during the LIA in many areas throughout the northern hemisphere (Barclay

*et al.*, 2009; Briner *et al.*, 2009; Menounos *et al.*, 2009). On the other hand, that some northern hemisphere glaciers exceeded or became very close to their late Holocene maxima prior to the LIA is not exceptional, and has been documented at a number of locations (e.g. Briner *et al.*, 2009; Maurer *et al.*, 2012; Schimmelfennig *et al.*, 2012).

The relatively minor expansion of Brooks Range glaciers during the LIA compared with expansions during the Neoglacial could be due to a progressive decrease in moisture availability through the late Holocene as Arctic sea ice expanded (England *et al.*, 2008; Antoniades *et al.*, 2011; Funder *et al.*, 2011). Sikorski *et al.* (2009) concluded that limited lowering of equilibrium line altitudes in the Brooks Range compared with elsewhere in Alaska during the LIA was due to drier-than-present conditions, consistent with the notion of moisture starvation due to expanded Arctic sea ice. This decrease in moisture may have been most pronounced in the Arctic; although middle Holocene advances are recognized elsewhere (Barclay *et al.*, 2009; Menounos *et al.*, 2009), to our knowledge the Brooks Range is the only location in the northern hemisphere yet identified where multiple glaciers seem to have expanded beyond their LIA positions as early as the middle Holocene.

## Conclusion

Our  $^{10}\text{Be}$  ages provide new evidence for the timing of deglaciation and Neoglaciation in the north-central Brooks Range. The  $^{10}\text{Be}$  inventory in bedrock surfaces in Roche Moutonnée valley appears to be significantly influenced by isotopic inheritance, which we attribute to the erosion-resistant beds within the Kanayut Conglomerate, possibly to weakly erosive polythermal glaciers. A  $^{10}\text{Be}$  age immediately outboard of late Holocene moraines in the upper Kurupa River valley (Fig. 2), supported by additional  $^{10}\text{Be}$  ages slightly downvalley, suggests that glaciers in the Brooks Range were either absent or smaller during the YD than during the Neoglacial. The lack of glacial advance associated with the YD agrees with other records in the Arctic, despite climate records showing pronounced YD cooling, including permafrost records that indicate significant winter cooling along the North Slope (Meyer *et al.*, 2010). It has been suggested that cooling during the YD in the Arctic was marked by strong seasonality with cold winters and increased sea-ice cover (Denton *et al.*, 2005) as well as aridity (Alley *et al.*, 1993), but with relatively mild summer conditions. In addition, glaciers in the Brooks Range are probably precipitation-sensitive (Sikorski *et al.*, 2009), and thus it is possible that strong aridity contributed to the lack of any extensive glacier expansion during the YD.

The  $^{10}\text{Be}$  ages for a middle Holocene moraine dating to ~4.6 ka provide the first numerical ages for the maximum Neoglacial expansion following early Holocene warmth (e.g. Kaufman *et al.*, 2004). Our  $^{10}\text{Be}$  ages agree with lichenometric ages of middle Holocene moraines within the 20% uncertainty of lichenometry based on the original linearly extrapolated growth curve of Ellis and Calkin (1984). This suggests that the lichenometrically dated moraine sequence, which features dozens of pre-LIA moraines across the Brooks Range, is generally robust. The  $^{10}\text{Be}$  ages confirm that glaciers in the Brooks Range reached similar extents during several glacier expansions during the Neoglacial. The effect of orbitally driven decreasing summer insolation in the Brooks Range may have been surpassed by decreasing moisture availability.

**Acknowledgements.** We thank Brandon Boldt and Samuel Kelley for assistance in the field and Nicolás Young for lab assistance. We would also like to thank Tom Hamilton for helpful discussions. Thanks to Fred

Luiszer for ICP measurements and Peter Bush for SEM assistance. Beata Csatho and Christopher Lowry reviewed an earlier version of this manuscript, and we think Brian Menounos and an anonymous reviewer for suggestions that greatly improved this manuscript. This work was supported by the National Science Foundation grant ARC-0902797 to J.P.B. and a graduate student research grant to M.E.B. from the Geological Society of America.

**Abbreviations.** LIA, Little Ice Age; YD, Younger Dryas.

## Supporting information

Additional supporting information can be found in the online version of this article:

**Supporting Figure S1.** Aerial photograph (same as in Figure 5) and lichen map from Lamb (1984) of Fireweed West Glacier. Bottom panel shows oblique photograph taken from airplane flight in July 2010 showing Fireweed West Glacier moraines.

**Supporting Figure S2.** Surficial geomorphic map from Lamb (1984) of the upper Kurupa River valley. See Figure 3 for aerial photograph of same area.

**Supporting Figure S3.** Aerial photograph and lichen map from Ellis and Calkin (1984) of the Triple East Glacier.

Please note: This supporting information is supplied by the authors, and may be re-organised for online delivery, but is not copy-edited or typeset by Wiley-Blackwell. Technical support issues arising from supporting information (other than missing files) should be addressed to the authors.

## References

- Alley RB, Meese DA, Shuman CA, *et al.* 1993. Abrupt increase in Greenland snow accumulation at the end of the Younger Dryas event. *Nature* **362**: 527–529.
- Antoniades D, Francus P, Pienitz R, *et al.* 2011. Holocene dynamics of the Arctic's largest ice shelf. *Proceedings of the National Academy of Sciences* **108**: 18,899–18,904.
- Applegate PJ, Urban NM, Keller K, *et al.* 2012. Improved moraine age interpretations through explicit matching of geomorphic process models to cosmogenic nuclide measurements from single landforms. *Quaternary Research* **77**: 293–304.
- Balascio NL, Kaufman DS, Briner JP, *et al.* 2005. Late Pleistocene glacial geology of the Okpilak-Kongakut Rivers region, northeastern Brooks Range, Alaska. *Arctic, Antarctic, and Alpine Research* **37**: 416–424.
- Balco G, Stone JO, Lifton NA, *et al.* 2008. A complete and easily accessible means of calculating surface exposure ages or erosion rates from  $^{10}\text{Be}$  and  $^{26}\text{Al}$  measurements. *Quaternary Geochronology* **3**: 174–195.
- Balco G, Briner JP, Finkel RC, *et al.* 2009. Regional beryllium-10 production rate calibration for late-glacial northeastern North America. *Quaternary Geochronology* **4**: 93–107.
- Barclay DJ, Wiles GC, Calkin PE. 2009. Holocene glacier fluctuations in Alaska. *Quaternary Science Reviews* **28**: 2034–2048.
- Berger A, Loutre MF. 1991. Insolation values for the climate of the last 10 million years. *Quaternary Science Reviews* **10**: 297–317.
- Briner JP, Kaufman DS. 2008. Late Pleistocene mountain glaciation in Alaska: key chronologies. *Journal of Quaternary Science* **23**: 659–670.
- Briner JP, Kaufman DS, Manley WF, *et al.* 2005. Cosmogenic exposure dating of late Pleistocene moraine stabilization in Alaska. *Geological Society of America Bulletin* **117**: 1108–1120.
- Briner JP, Davis PT, Miller GH. 2009. Latest Pleistocene and Holocene glaciation of Baffin Island, Arctic Canada: key patterns and chronologies. *Quaternary Science Reviews* **28**: 2075–2087.
- Briner JP, Young NE, Goehring BM, *et al.* 2012. Constraining Holocene  $^{10}\text{Be}$  production rates in Greenland. *Journal of Quaternary Science* **27**: 2–6.
- Brogé WP, Reiser HN, Dutro JT, *et al.* 1979. Bedrock geologic map of the Philip Smith Mountains Quadrangle, Alaska. *U.S. Geologic Survey, Miscellaneous Field Studies Map Scale* **1**: 250,000.

- Calkin PE, Ellis JM. 1980. A lichenometric dating curve and its application to Holocene glacier studies in the central Brooks Range, Alaska. *Arctic and Alpine Research* **12**: 245–264.
- Denton GH, Alley RB, Comer GC, *et al.* 2005. The role of seasonality in abrupt climate change. *Quaternary Science Reviews* **24**: 1159–1182.
- Denton GH, Anderson RF, Toggweiler JR, *et al.* 2010. The last glacial termination. *Science* **328**: 1652–1656.
- Ellis JM. 1982. *Holocene glaciation of the Central Brooks Range, Alaska*. PhD Dissertation, University at Buffalo.
- Ellis JM, Calkin PE. 1984. Chronology of Holocene glaciation, Central Brooks Range, Alaska. *Bulletin of the Geological Society of America* **95**: 897–912.
- England JH, Lakeman TR, Lemmen DS, *et al.* 2008. A millennial-scale record of Arctic Ocean sea ice variability and the demise of the Ellesmere Island ice shelves. *Geophysical Research Letters* **35**: L19502.
- Fenton CR, Hermanns RL, Blikra LH, *et al.* 2011. Regional  $^{10}\text{Be}$  production rate calibration for the past 12 ka deduced from the radiocarbon-dated Grtlandsura and Russenes rock avalanches at 69N, Norway. *Quaternary Geochronology* **6**: 437–452.
- Fisher D, Zheng J, Burgess D, *et al.* 2012. Recent melt rates of Canadian arctic ice caps are the highest in four millennia. *Global and Planetary Change* **84–85**: 3–7.
- Funder S, Goosse H, Jepsen H, *et al.* 2011. A 10,000-year record of Arctic Ocean sea-ice variability-view from the beach. *Science* **333**: 747–750.
- Gardner AS, Moholdt G, Wouters B, *et al.* 2011. Sharply increased mass loss from glaciers and ice caps in the Canadian Arctic Archipelago. *Nature* **473**: 357–360.
- Gosse JC, Evenson EB, Klein J, *et al.* 1995. Precise cosmogenic  $^{10}\text{Be}$  measurements in western North America; support for a global Younger Dryas cooling event. *Geology* **23**: 877–880.
- Hamilton TD. 1982. Quaternary stratigraphic sections with radiocarbon dates, Killik River Quadrangle, Alaska. *U.S.G.S. Open File Report*: 82–606.
- Hamilton TD. 1986. Late Cenozoic glaciation of the Central Brooks Range. In: Hamilton TD, Reed KM, Thorson RM (Eds). *Glaciation in Alaska-The geologic record*. Alaska Geological Society, Anchorage; pp. 9–50.
- Hamilton TD. 2003. Surficial geology of the Dalton Highway (Itkillik-Sagavanirk Rivers) area, southern Arctic Foothills, Alaska. *Professional Report-Alaska Division of Geological and Geophysical Surveys* 32.
- Harrison S, Glasser N, Anderson E, *et al.* 2010. Late Pleistocene mountain glacier response to North Atlantic climate change in southwest Ireland. *Quaternary Science Reviews* **29**: 3948–3955.
- Heyman J, Stroeven AP, Harbor JM, *et al.* 2010. Too young or too old: evaluating cosmogenic exposure dating based on an analysis of compiled boulder exposure ages. *Earth and Planetary Science Letters* **302**: 71–80.
- Holland MM, Bitz CM. 2003. Polar amplification of climate change in coupled models. *Climate Dynamics* **21**: 221–232.
- Ingólfsson Ó, Björck S, Hafliðason H, *et al.* 1997. Glacial and climatic events in Iceland reflecting regional North Atlantic climatic shifts during the Pleistocene–Holocene transition. *Quaternary Science Reviews* **16**: 1135–1144.
- Jacob T, Wahr J, Pfeffer WT, *et al.* 2012. Recent contributions of glaciers and ice caps to sea level rise. *Nature* **482**: 514–518.
- Kaufman DS, Ager TA, Anderson NJ, *et al.* 2004. Holocene thermal maximum in the western Arctic (0–180W). *Quaternary Science Reviews* **23**: 529–560.
- Kaufman DS, Anderson RS, Hu FS, *et al.* 2010. Evidence for a variable and wet Younger Dryas in southern Alaska. *Quaternary Science Reviews* **29**: 1445–1452.
- Kelly MA, Lowell TV, Hall BL, *et al.* 2008. A  $^{10}\text{Be}$  chronology of late glacial and Holocene mountain glaciation in the Scoresby Sund region, east Greenland: implications for seasonality during late glacial time. *Quaternary Science Reviews* **27**: 2273–2282.
- Kokorowski HD, Anderson PM, Mock CJ, *et al.* 2008. A re-evaluation and spatial analysis of evidence for a Younger Dryas climatic reversal in Beringia. *Quaternary Science Reviews* **27**: 1710–1722.
- Lal D. 1991. Cosmic ray labeling of erosion surfaces: in situ nuclide production rates and erosion models. *Earth and Planetary Science Letters* **104**: 424–439.
- Lamb B. 1984. *Geomorphology of the upper Kurupa River valley, Brooks Range, Alaska*. MS Thesis, University at Buffalo.
- Lohne ØS, Mangerud J, Svendsen JJ. 2012. Timing of the Younger Dryas glacial maximum in western Norway. *Journal of Quaternary Science* **27**: 81–88.
- Mangerud J, Landvik JY. 2007. Younger Dryas cirque glaciers in western Spitsbergen; smaller than during the Little Ice Age. *Boreas* **36**: 278–285.
- Mann DH, Peteet DM, Reanier RE, *et al.* 2002. Responses of an arctic landscape to Lateglacial and early Holocene climatic changes: the importance of moisture. *Quaternary Science Reviews* **21**: 997–1021.
- Maurer MK, Menounos B, Luckman H, *et al.* 2012. Late Holocene glacier expansion in the Cariboo and northern Rocky Mountains, British Columbia, Canada. *Quaternary Science Reviews* **19**: 71–80.
- Meier MF, Dyurgerov MB, McCabe GJ. 2003. The health of glaciers; recent changes in glacier regime. *Climate Change* **59**: 123–135.
- Menounos B, Osborn G, Clague JJ, *et al.* 2009. Latest Pleistocene and Holocene glacier fluctuations in western Canada. *Quaternary Science Reviews* **28**: 2049–2074.
- Meyer H, Schirmermeister L, Yoshikawa K, *et al.* 2010. Permafrost evidence for severe winter cooling during the Younger Dryas in northern Alaska. *Geophysical Research Letters* **37**: L03501.
- Miller GG, Brigham-Grette J, Alley RB, *et al.* 2010. Temperature and precipitation history of the Arctic. *Quaternary Science Reviews* **29**: 1679–1715.
- Molina BF. 2007. Late nineteenth to early twenty-first century behavior of Alaskan glaciers as indicators of changing regional climate. *Global and Planetary Change* **56**: 23–56.
- Nishiizumi K, Imamura M, Caffè MW, *et al.* 2007. Absolute calibration of  $^{10}\text{Be}$  AMS standards. *Nuclear Methods and Instruments in Physics Research B* **258**: 403–413.
- Overpeck J, Hughen K, Hardy D, *et al.* 1997. Arctic environmental change of the last four centuries. *Science* **278**: 1251–1256.
- Porter SC. 2007. Neoglaciation in the American Cordilleras. In: Elias S (Ed.). *Encyclopedia of Quaternary Science*. Elsevier: Amsterdam; 1133–1142.
- Schimmelpennig I, Schaefer JM, Akçar N, *et al.* 2012. Holocene glacier culminations in the Western Alps and their hemispheric relevance. *Geology* **40**: 891–894.
- Serreze MC, Barry RG. 2011. Processes and impacts of Arctic amplification: a research synthesis. *Global and Planetary Change* **77**: 85–96.
- Sharp M, Burgess DO, Cogley JG, *et al.* 2011. Extreme melt on Canada's Arctic ice caps in the 21<sup>st</sup> century. *Geophysical Research Letters* **38**: L11501.
- Sikorski JJ, Kaufman DS, Manley WF, *et al.* 2009. Glacial–geologic evidence for decreased precipitation during the Little Ice Age in the Brooks Range, Alaska. *Arctic, Antarctic, and Alpine Research* **41**: 138–150.
- Solomina O, Calkin PE. 2003. Lichenometry as applied to moraines in Alaska, USA, and Kamchatka, Russia. *Arctic, Antarctic, and Alpine Research* **35**: 129–143.
- Stone JO. 2000. Air pressure and cosmogenic isotope production. *Journal of Geophysical Research* **105**: 23753–23759.
- Stuiver M, Grootes PM. 2000. GISP2 Oxygen isotope ratios. *Quaternary Research* **53**: 277–284.