



Abrupt warming and alpine glacial retreat through the last deglaciation in Alaska interrupted by modest Northern Hemisphere cooling

Joseph P. Tulenko^{1,2}, Jason P. Briner², Nicolás E. Young³, and Joerg M. Schaefer³

¹Berkeley Geochronology Center, 2455 Ridge Road, Berkeley, CA 94709, USA

²Department of Geology, University at Buffalo, 126 Cooke Hall, Buffalo, NY 14260, USA

³Lamont-Doherty Earth Observatory, Columbia University, 61 Route 9W, Palisades, NY 10964, USA

Correspondence: Joseph P. Tulenko (jtulenko@bgc.org)

Received: 14 September 2023 – Discussion started: 28 September 2023

Revised: 29 January 2024 – Accepted: 3 February 2024 – Published: 21 March 2024

Abstract. Alpine glacier-based temperature reconstructions spanning the last deglaciation provide critical constraints on local to regional climate change and have been reported from several formerly glaciated regions around the world yet remain sparse from high-northern-latitude regions. Using newly and previously ¹⁰Be-dated moraines, we report paleo-glacier equilibrium line altitudes (ELAs) for 15 time slices spanning the Last Glacial Maximum (LGM) to the Little Ice Age (LIA) for a valley in the western Alaska Range. We translate our ELA reconstructions into a proxy for summer temperature by applying a dry adiabatic lapse rate at each reconstructed ELA relative to the outermost LIA moraine. We observe ~ 4 °C warming through the last deglaciation at our site that took place in two steps following initial gradual warming: ~ 1.5 °C abrupt warming at 16 ka, ~ 2 kyr after the onset of global CO₂ rise, and ~ 2 °C warming at ~ 15 ka, near the start of the Bølling. Moraine deposition and modest summer cooling during Heinrich Stadial 1 and the early Younger Dryas (YD) suggest that despite these events being expressed more strongly in wintertime, the classic blueprint of North Atlantic climate variability extends to the western Arctic region.

temperature changes on both centennial (Roe et al., 2017) and millennial timescales (Schaefer et al., 2006). The last deglaciation (~ 19–11 ka) contains periods of abrupt warming akin to contemporary climate change – particularly later in the interval during the “late glacial” at ~ 15–11 ka – and considerable effort has been spent constraining the timing and pace of mountain glacier fluctuations worldwide through this interval (e.g., Putnam et al., 2013; Ivy-Ochs, 2015; Palascios et al., 2020). However, mountain glacier fluctuations in the high northern latitudes – in locations where ice sheets were largely absent or restricted – are less well characterized due in part to the limited availability of suitable sites for organic radiocarbon dating, which presents a significant challenge compared to more temperate regions (Wittmeier et al. 2020).

Glacier equilibrium line altitudes (ELAs) are a representative measure of the climatic conditions supporting a glacier. Climate-driven positive and negative mass balance changes force the ELA to lower or rise, respectively (Ohmura et al., 1992; Oerlemans et al., 2005; Braithwaite, 2008). Reconstructing paleo-glaciers through detailed mapping and dating techniques provides valuable data for estimating paleo-ELAs, which serve as a powerful paleoclimate proxy for glaciated regions. Improved access to accurate digital elevation models (DEMs) and the development of semiautomated paleo-glacier surface and ELA reconstruction tools (Pellitero et al., 2015, 2016) pave a path for more objective measures of paleo-ELA calculation. Changes in ELA are dependent primarily on summer (JJA) temperatures and annual snow-

1 Introduction

Global mountain glacier recession is one of the clearest indicators of modern planetary warming (Hugonnet et al., 2021). Paleo-data suggest that alpine glaciers are highly sensitive to




Figure 1. (a) Moraine map of the North Swift River valley with reconstructed moraine positions denoted with red lines ($n = 15$). Color overlays denote broad morphostratigraphic equivalence of landforms and are arranged as darker shades being stratigraphically older than lighter shades. Blue shades are MIS-4-age moraines, brown-red shades into yellow shades are LGM and deglacial moraines, and green are Holocene-age moraines. Inset: starred location of the Revelation Mountains in Alaska. (b) Schematic of alpine glacier retreat in the North Swift River valley that includes moraine ages reported here and in Tulenko et al. (2022). Panel (a) source: ArcticDEM product available from the University of Minnesota Polar Geospatial Center. Inset map source: IBCAO product available from GEBCO.

fall (Oerlemans et al., 2005). Some studies suggest, however, that in most environments other than the most extremely arid environments such as modern-day inner Mongolia, alpine glacier ELAs are dominated by summer temperatures (e.g., Rupper and Roe, 2008).

Building on a recently published ^{10}Be chronology of 14 moraines in the North Swift River valley, Revelation Mountains, western Alaska Range (Tulenko et al., 2018, 2022), we constrain additional ^{10}Be ages for three late Holocene moraines in the valley and generate a paleo-ELA-based proxy summer temperature record from our site. We estimate the timing and pace of warming through the last deglaciation by transforming ΔELAs into summer temperature assuming a dry adiabatic lapse rate of 10°C km^{-1} . The magnitude of temperature change we obtain is supported by independent climate proxy records in Alaska, and we compare our record with other glacier-based temperature reconstructions

to speculate about the controls on climate in Alaska through deglaciation.

2 The Revelation Mountains field site

The Revelation Mountains field site is located on the western limb of the Alaska Range (Fig. 1 inset) where dense sequences of moraines deposited by alpine glaciers fluctuating through the last deglaciation are well preserved. The Revelation Mountains are cored by a granitic pluton, and large tabular boulders deposited on moraines are suitable targets for cosmogenic ^{10}Be exposure dating. The site is located far south from seismic activity along the Denali Fault, and thus moraines deposited in these valleys have likely remained more stable after deposition compared to moraines in locations more proximal to the Denali Fault system (e.g., Briner et al., 2005; Matmon et al., 2010; Dortch et al., 2010). All these factors have led to productive chronologic constraints on deglaciation in Alaska (Tulenko et al., 2022).

Within the Revelation Mountains, the North Swift River valley hosts the densest and most comprehensive sequence of moraines of any valley surveyed at our site (Fig. 1). For more details on the mapping and dating approaches used to generate the deglacial chronology in the North Swift River valley site, readers are referred to Tulenko et al. (2022). After a thorough moraine dating campaign of deglacial moraines in the valley, we additionally mapped and dated moraines in the valley deposited near the terminus of the extant glacier that were likely late Holocene in age (Fig. 2). For this study, the purpose was to target late Holocene moraines to contextualize the magnitude of deglacial retreat in the North Swift River valley. Thus, all inferred climate fluctuations reported here are relative to the late Holocene advances observed at our site.

3 Cosmogenic ^{10}Be exposure dating of late Holocene glacier fluctuations in the Revelation Mountains

To complete the time series of glacier retreat in the North Swift River valley, we collected surface samples in summer 2019 for ^{10}Be dating from 15 granitic boulders on three distinct moraine crests between the innermost late glacial moraine (F2-IV E) and an unnamed extant glacier (Figs. 1 and 2). The moraines are sharp-crested and clast-supported with minimal vegetation cover (Fig. 3). We selected the largest tabular boulders ($> 1\text{--}2\text{ m}$ tall) with horizontal to sub-horizontal top surfaces to avoid excessive snow cover. Surface samples approximately 2 cm thick (see Table 1 for exact thickness measurements) and $\sim 1\text{ kg}$ in mass were extracted using a Hilti brand angle grinder with diamond-tipped cutting disks and a hammer and chisel. GPS coordinates were taken with a handheld GPS, and shielding corrections were calculated using in-field measurements with a handheld clinome-




Figure 2. (a) Shaded relief map of the modern North Swift River valley glacier and late Holocene moraines with ^{10}Be ages displayed. Samples listed in italics are suspected outliers. The age for the F2-IV E moraine segment is from Tulenko et al. (2022). Map source: 5 m resolution IFSAR Alaska DEM product available from the USGS. (b) Normal probability density functions (PDFs) for individual ages are in red and summed PDFs for the two clusters of ages around 750 and 320 a are in black. Ages older than expected and well beyond 2 standard deviations from other ages in the dataset are in gray.

ter and the shielding correction calculator from the online exposure age calculator website (<https://hess.ess.washington.edu/>, last access: 1 September 2023). We targeted large, tabular boulders on clast-supported moraines to minimize the likelihood that post-depositional processes – e.g., snow cover and moraine degradation – inhibited production of cosmogenic ^{10}Be in sampled surfaces, which would produce ages younger than the true timing of moraine deposition.

All samples were processed from full surface sample to prepared ^{10}Be targets at the Lamont-Doherty Earth Observatory (LDEO) cosmogenic dating laboratory following procedures outlined in Schaefer et al. (2009). Beryllium measurements for samples and batch blanks were made at the Lawrence Livermore National Lab Center for Accelerator Mass Spectrometry (LLNL-CAMS). All $^{10}\text{Be}/^9\text{Be}$ ratios were standardized to the 07KNSTD standard of 2.85×10^{-12} (Nishiizumi et al., 2007), and blank-corrected final ^{10}Be concentrations are reported in Table 1 and in the ICE-D database (<https://www.ice-d.org>, last access: 1 September 2023) along with all other relevant sample observations and measurements.

^{10}Be surface exposure ages reported here were calculated on the online exposure age calculator (Balco et al., 2008) using the Baffin Bay Arctic production rate (Young et al., 2013) and the time-dependent scaling scheme from Lal (1991) and Stone (2000). Several previous studies in Alaska have utilized this production rate calibration–scaling scheme combination (Valentino et al., 2021; Tulenko et al., 2022, and references cited therein), and the production rate measured in Baffin Bay is statistically identical to other high-latitude Northern Hemisphere calibration sites such as the Scottish Highlands Rannoch Moor site (Putnam et al., 2019), the Swiss Alps Chironico Landslide site (Claude et al., 2014), and northeastern North America (Balco et al., 2009). Use of other commonly applied production rate calibrations would result in ages offset by $\sim 3\%$. We do not attempt to correct for snow cover. We also do not correct for boulder surface erosion since the boulders are crystalline granitic lithology and thus resistant to weathering, particularly in the brief (hundreds of years) period between deposition and sample measurement. Ages from this study converted to common era (CE); below) use 2019 (date of collection) as the reference starting date.

Ages from the three dated moraines situated 200–1000 m inboard of the innermost late Pleistocene (F2-IV E) moraine range from 791 ± 20 – 304 ± 15 a (1228–1715 CE ($n = 10$; not including five potential outliers; Figs. 1, 2, Table 1). Ages that are suspected outliers do not overlap with any other ages within 2 standard deviations and are older than the remaining ages (Fig. 2b), which indicates they may have been influenced by nuclide inheritance. The outermost moraine (LH-1) has four ages that overlap within 2 standard deviations and average 755 ± 35 a (1263 ± 35 CE). The next moraine (LH-2) does not have any ages that overlap, but one boulder age at 737 ± 15 a (1282 ± 15 CE) coincides with the ages from LH-1 so it is possible it was deposited inboard of the LH-1 position at the same time the LH-1 moraine formed and was then recycled during the advance that formed LH-2. For the innermost moraine dated, two of the five boulder ages overlap within 2 standard deviations and average 323 ± 19 a (1696 ± 19 CE). Additionally, LH-3 contains one boulder age that also conforms well to the ages from LH-1 at 721 ± 20 a (1298 ± 20 CE) and could have a similar history to the age from the LH-2 moraine at 737 ± 15 a (1282 ± 15 CE).

Many of the well-constrained observations of Little Ice Age (LIA) glacier fluctuations in Alaska come from southern Alaska, where tree-ring chronologies from trees sheared at their bases by glacial overriding tightly constrain episodes of glacier advance (Figs. 4 and 5; Wiles et al., 2002, 2004; Barclay et al., 2009). These precise chronologies reveal two major glacier advances at ~ 1200 – 1300 CE and ~ 1600 – 1800 CE (Fig. 5). In comparison, we observe moraines deposited in the North Swift River valley at 755 ± 35 a (1263 ± 35 CE) and 323 ± 19 a (1696 ± 19 CE). The coherence of ^{10}Be ages in the North Swift River valley with tree-kill dates in southern Alaska suggests similar climate controls for glaciers

Table 1. ^{10}Be age information.

Sample name	Latitude (DD)	Longitude (DD)	Elevation (m a.s.l.)	Thickness (cm)	Shielding correction	Qtz dissolved (g)	Carrier added (g)	[Be-10] / [Be-9] (10^{-14})	[Be-10] Atoms g^{-1}	Age (a)	Age (CE)
LH-1											
19AK-75 ^a	61.67542	-154.28345	822	1.85	0.964221	53	0.1815	3.013 ± 0.092	7106 ± 217	787 ± 24	1232 ± 24
19AK-76 ^a	61.67548	-154.28334	822	2.41	0.964221	48	0.182	4.661 ± 0.099	12194 ± 259	1362 ± 29	657 ± 29
19AK-77 ^a	61.67552	-154.28322	818	4.92	0.964221	70	0.1822	3.509 ± 0.106	6259 ± 189	713 ± 22	1306 ± 22
19AK-78 ^a	61.67542	-154.28313	820	2.52	0.964221	70	0.1819	3.637 ± 0.101	6521 ± 182	727 ± 20	1292 ± 20
19AK-79 ^a	61.67546	-154.28278	820	2.87	0.964221	62	0.1821	3.526 ± 0.087	7071 ± 175	791 ± 20	1228 ± 20
LH-2											
19AK-70 ^a	61.6741	-154.28181	829	2.5	0.964221	70	0.1806	3.761 ± 0.081	6658 ± 143	737 ± 16	1282 ± 16
19AK-71 ^b	61.6739	-154.28194	829	2.9	0.964221	71	0.1814	28.607 ± 0.532	50099 ± 931	5591 ± 104	-3572 ± 104
19AK-72 ^b	61.67412	-154.28212	828	2.5	0.964221	70	0.182	16.567 ± 0.269	29637 ± 481	3312 ± 54	-1293 ± 54
19AK-73 ^a	61.6741	-154.28217	828	2.2	0.964221	70	0.1822	2.433 ± 0.074	4375 ± 133	484 ± 15	1535 ± 15
19AK-74 ^a	61.67403	-154.28236	832	2.67	0.964221	72	0.1807	4.940 ± 0.099	8563 ± 173	946 ± 19	1073 ± 19
LH-3											
19AK-65 ^b	61.67059	-154.27878	864	2.77	0.964245	71	0.1813	11.047 ± 0.206	19445 ± 362	2099 ± 39	-80 ± 39
19AK-66 ^b	61.67073	-154.27852	863	2.83	0.964245	69	0.1812	1.568 ± 0.077	2830 ± 140	304 ± 15	1715 ± 15
19AK-67 ^b	61.67073	-154.27855	862	2.04	0.964245	69	0.1822	3.715 ± 0.104	6729 ± 188	721 ± 20	1298 ± 20
19AK-68 ^b	61.67081	-154.27794	862	2.82	0.964245	71	0.1823	3.182 ± 0.082	5635 ± 145	608 ± 16	1411 ± 16
19AK-69 ^b	61.67096	-154.27766	865	2.38	0.964245	73	0.1816	1.869 ± 0.065	3193 ± 112	342 ± 12	1677 ± 12

^a The blank ratio for these samples is 7.685×10^{-16} . ^b The blank ratio for these samples is 5.311×10^{-16} . Carrier concentration for all samples and blanks was 1029.6 ppm. Rock density is assumed to be 2.65 g cm^{-3} for all samples. Zero surface erosion assumed. Ages presented here are calculated using the Arctic production rate (Young et al., 2013) and time-dependent scaling from Lal (1991) and Stone (2000).



Figure 3. Example photos of sampled boulders on the three late Holocene moraines (top and bottom left panels). The bottom right panel is an image of the left-lateral segment of LH-3 where sampled boulders are situated. Note the person on the moraine crest for scale. Images of all sampled boulders can be found in the ICE-D database (<https://www.ice-d.org>, last access: 1 September 2023).




Figure 4. Locations of glaciers in southern Alaska with well-constrained late Holocene advances from tree-kill dates. The location of the Revelation Mountains is denoted by a star in the inset. Base map available from ASTER.

in the western Alaska Range and in southern Alaska during the LIA.

While the sampling techniques discussed above greatly reduce the likelihood of post-depositional processes impacting age determinations, it is virtually impossible to detect in the field which samples may have excessive cosmogenic nuclide inventories (i.e., nuclide inheritance) from surface exposure prior to glacier advances and moraine-building events. Commonly with erosive (non-cold-based) alpine glaciers, measured samples on moraines impacted by nuclide inheritance manifest as extreme values that do not conform to any other ages in the distribution (e.g., Heyman et al., 2011). Thus, we suspect all that samples observed on our three moraines with extreme old values that do not conform to any other ages were impacted by nuclide inheritance. Not considering the older extreme values across the three moraines ($n = 5$), we find 8 of the remaining 10 samples overlapping in two clusters, ~ 1200 – 1300 CE ($n = 6$) and ~ 1700 CE ($n = 2$). We tentatively conclude that because these samples are both internally consistent and consistent with other independent records of LIA glacier fluctuations in Alaska, they may reflect the true age of two LIA glacier advances at our site.

4 Glacier surfaces, ELAs, and climate reconstructions in the Revelation Mountains through deglaciation

To construct a climate record built on the completed chronology at our site, we employ the tools developed for use in ArcGIS from Pelitiero et al. (2015, 2016) to first reconstruct paleo-glacier surfaces and then equilibrium line altitudes (ELAs) for each of the previously dated moraine positions ($n = 14$) and the outermost late Holocene moraine position (LH-1; 15 total moraines). We select the outermost late Holocene moraine because it is the most robustly dated and the late Holocene moraines are all within 1 km of each other, resulting in similar paleo-ELA values. The glacier surface reconstruction tool (GLARE; Pelitiero et al., 2016) is semiautomated but requires the following inputs: a base map digital elevation model (DEM), polygons of maximum glacier outline constraints, and glacier center flowline polygons (Fig. 6). We mapped glacier outlines on a 5 m resolution IFSAR DEM base map provided by the USGS and available for download from National Map (<https://apps.nationalmap.gov/downloader/>, last access: 20 August 2023). Maximum constraints for each individual paleo-glacier position were outlined based on terminal and lateral moraines originally mapped in Tulenko et al. (2018) and extending up to the




Figure 5. Comparison of late Holocene moraine ages (green dots; gray dots are outliers) with tree-ring-based southern Alaska glacial fluctuations (black bars; Barclay et al., 2009). Light blue vertical bars indicate periods of moraine deposition in the Revelation Mountains (upper panel) and tree-kill dates signifying glacier culminations in southern Alaska (lower panels) at ~ 1200 – 1300 CE and ~ 1600 – 1800 CE. See Fig. 4 for geographic locations of tree-kill dates.

valley drainage divide in the steeper terrain once moraines were no longer visible. In steeper terrain, it is difficult to find preserved evidence of glacier extents, so the divide offers the maximum possible limit of glacial extent. For the remaining steps, we resampled the 5 m resolution DEM down to 100 m resolution to smooth out the surface. We suggest that since post-glacial landscape evolution has likely altered the land surface at fine scale, a smoothed surface may minimize any influence of post-glacial landscape evolution on our glacier surface reconstructions. Using the resampled DEM and the series of watershed analysis tools in ArcGIS, we produced ice flowline polylines that generally follow the modern drainages (Fig. 6). The GLARE tool then calculates ice thickness along the center flowline assuming perfect plasticity for ice rheology and a value for basal shear stress that we set to 100 kPa for all paleo-glaciers at all positions along their center flowlines. Finally, we use the topo-to-raster interpolation method (see Hutchinson et al., 2011, for details) supplied by the GLARE tool to reconstruct a glacier surface extending outward from center flowline thickness values of each paleo-glacier and confined either by their respective glacier outline where moraines exist or the valley wall in steeper terrain (see Fig. 6 insets for examples).

Following glacier surface reconstructions, we employ the accumulation area ratio method supplied by the ArcGIS ELA

Table 2. Glacier equilibrium line altitude information.

Moraine name	Mean ELA (m)	ELA error (m)	Moraine age (ka)	Moraine age error (ka)
F2-I A	975	16	21.3	0.8
F2-I B	1019	18	20.1	1.3
F2-II A	1028	14	19.5	1.0
F2-II B	1030	14	18.7	0.8
F2-III A	1031	13	16.9	1.2
F2-III B	1043	18	16.0	0.9
F2-III C	1072	20	15.9	0.9
F2-III D	1141	22	16.6	0.6
F2-III E	1152	23	15.7	0.7
F2-IV A	1172	23	16.1	0.9
F2-IV B	1175.5	23.5	15.0	0.7
F2-IV C	1182.5	23.5	15.5	0.7
F2-IV D	1206	26	15.7	0.7
F2-IV E	1362.5	20.5	12.8	0.6
LH-1	1378.5	20.5	0.75	0.05

Notes: all ELAs are measured using 0.63 ± 0.04 accumulation area ratios. All ages reported here are calculated in Tulenko et al. (2022).

tool (Pelitero et al., 2015) for each paleo-glacier. We assume a ratio of 0.63 ± 0.04 , which has been suggested as representative for alpine glaciers greater than 4 km^2 (Kern and Laszlo, 2010). ELA values for each glacier position, along with age constraints reported in Tulenko et al. (2022), can be found in Table 2. We chose the AAR method as opposed to the accumulation area balance ratio (AABR) method, even though it is likely that the AABR method better approximates modern glacier ELAs (Pelitero et al., 2015) for two reasons. First, there are required observations for the AABR method that are difficult to quantify for paleo-glaciers, namely the balance ratio, which must be linear for the method to be valid (Osmaston, 2005). Second, since we apply the same exact methodologies for each paleo-glacier reconstructed, we assume that any changes to methodologies (e.g., here using AAR vs. AABR, using different basal shear stress values) would result in systematic shifts that would not appreciably impact the magnitude of ELA change observed. As a final observation, a recent study of paleo-glacier and ELA reconstructions Alaska-wide used both AAR and AABR methods for calculating ELAs and found minimal difference in their results (Walcott et al., 2024). All points considered, we elected to use the simpler, but still effective, AAR method.

Finally, we calculate summer temperature anomalies for each glacier position relative to the LIA by applying the dry adiabatic lapse rate of $10 \text{ }^\circ\text{C km}^{-1}$ to each ΔELA . The dry adiabatic lapse rate is applied because it serves as an end-member; over timescales greater than a few hours, it is not possible for an air mass to sustainably exceed the dry adiabatic lapse rate. Thus, our reconstruction likely provides a maximum estimate in the magnitude of temperature change




Figure 6. Plan view of the North Swift River valley and reconstructed glacier extents in the main panel. Outlines of every glacier surface and extent reconstructed in this analysis (in red). Blue lines are the flowlines reconstructed on the resampled, lower-resolution DEM. In the insets are 3D visualizations of reconstructed glacier surfaces. ELA reconstructions at each glacier position viewed in oblique 3D with $4\times$ vertical exaggeration. Each number in the main panel corresponds to the toe of the glacier position reconstructed in each inset panel. Glacier surfaces 1–6 are viewed from the perspective in the bottom left corner of the main panel, and glacier surface 7 is viewed from the perspective near the top right.

at each glacier position because lower lapse rates would translate to smaller changes in atmospheric temperature.

Our ELA reconstructions indicate ~ 400 m of total ELA rise through the deglacial interval (975–1362.5 m a.s.l.), which translates to approximately 4°C of warming (Fig. 7, Table 2). While there was notable lateral recession between 21 and 16 ka, the valley floor here is low-sloping, and we observe minimal ELA rise (~ 50 m) and modest warming ($\sim 0.4^\circ\text{C}$). Following this interval, a period of abrupt ELA rise corresponding to $\sim 1.5^\circ\text{C}$ of warming occurred, followed by a pause in deglaciation at ~ 16 – 15 ka when several moraines were deposited. After this period, a second interval of net $\sim 2^\circ\text{C}$ warming from ~ 15 to 12.8 ka led to glacier recession and non-deposition of moraines until the final late glacial moraine preserved in our valley was deposited at 12.8 ± 0.6 ka with an ELA of 1362.5 m a.s.l., only 16 m below the ELA of the outermost late Holocene moraine in the valley.

5 Discussion

Given that modern average lapse rates in Alaska may be as low as $\sim 4.3^\circ\text{C km}^{-1}$ (Verbyla and Kurkowski, 2019), temperature estimates through the last deglaciation could be less than half the magnitude reported here (2°C total warming through deglaciation). However, additional proxy evidence across Alaska spanning deglaciation from pollen assemblages (Viau et al., 2008), fossil chironomids (Kurek et al., 2009), and leaf wax hydrogen isotope data (Daniels et al., 2021) – as well as Gulf of Alaska sea surface temperatures (Praetorius et al., 2020) – suggests $\sim 4^\circ\text{C}$ summer temperature depressions during the LGM, consistent with the glacier-based temperature reconstruction from the Revelation Mountains. Several data assimilation products based on marine sediment records further substantiate the magnitude of cooling observed in our record (Tierney et al., 2020). Moreover, multiple lines of proxy evidence from Alaska indicate that conditions were drier than present through much of the last deglaciation until the late Younger Dryas (YD) (Viau et al., 2008; Finkenbinder et al., 2014, 2015; Dorfman et al., 2015), suggesting that the time-averaged late Pleistocene en-




Figure 7. Comparison of the Revelation Mountains retreat chronology with Southern Hemisphere and Alps glacial retreat, as well as global and regional climate forcing mechanisms. Top to bottom: the Revelation Mountains ELA-based temperature curve (this study), composite ELA reconstructions from the Alps (Ivy-Ochs, 2015), temperature curve from Ohau Glacier, NZ (Putnam et al., 2013), composite temperature curve from Patagonian glaciers (Denton et al., 1999; Strelin et al., 2011), composite atmospheric CO₂ concentrations from Antarctic ice cores (Bereiter et al., 2015), and Greenland-wide average mean annual temperature from ice cores (20–11 ka; Buizert et al., 2014, 11–0 ka; Kobashi et al., 2017).

environmental lapse rate was likely higher than modern and closer to the end-member dry lapse rate used here. However, there is a notable increase in precipitation observed following the YD in Alaska (e.g., Viau et al., 2008; Kaufman et al., 2010) that could have modulated subsequent glacier advances in the Holocene. Thus, this shift in precipitation may have enhanced the LIA advances at our site to nearly match the youngest late glacial moraine (F2-IV) despite a comparatively warmer Holocene climate. In this scenario, by not accounting for changes in precipitation, we would be underestimating either the magnitude of warming from the late glacial into the Holocene or the magnitude of late glacial cooling relative to the LIA. While we still lean on evidence suggesting that summer temperature is the most important climatic factor for alpine glaciers (Rupper and Roe, 2008), and the total magnitude of warming observed at our site using our end-member lapse rate (10 °C km^{-1}) is consistent with other lines of proxy evidence, further examination of the role of precipitation variations through deglaciation and the Holocene in

glacier advances at our site and others across Alaska is likely warranted.

Moraine ages from across Alaska reveal an emerging pattern, although in no other valley is the moraine record as complete or well dated as in the Revelation Mountains. The Last Glacial Maximum advance culminated in Alaska at $\sim 21\text{--}19\text{ ka}$ (Kaufman et al., 2011, Tulenko et al., 2018). Additionally, several moraines have been dated between ~ 16 and 15 ka during the later stage of Heinrich Stadial 1 (HS-1). In rare cases, most notably in the Revelation Mountains and in the Ahklun Mountains (e.g., Young et al., 2019), moraines date to $\sim 12.8\text{--}12.1\text{ ka}$ during the late glacial and within the earliest and coldest interval of the Younger Dryas (YD). Despite limited chronologic constraints, the magnitude of glacier retreat observed in other valleys in Alaska appears to track with glaciers in the Revelation Mountains (Tulenko et al., 2022). Thus, it is possible the last deglaciation climate reconstruction from the Revelation Mountains may be representative of the broader Alaska region.

Following the last deglaciation, we observe minimal glacier activity at our site through much of the Holocene until the LIA, when glacier advances nearly matched the extent of the youngest late glacial advance. While this is consistent with glacier behavior at several sites in Alaska, we note the contrasting pattern of Holocene glacier activity observed in valleys in the Brooks Range. Observations suggest that some neoglacial advances predating the LIA were slightly more extensive in the region (e.g., Ellis and Calkin, 1984; Badding et al., 2013; Pendleton et al., 2017). Moreover, in the Ahklun Mountains, Young et al. (2019) observe two minor glacier re-advances in the Waskey valley at 12.5 ± 0.1 and $12.1 \pm 0.4\text{ ka}$. While there appears to be no preserved evidence of glacier advances between the $12.8 \pm 0.6\text{ ka}$ moraine and the LIA moraines in the Revelation Mountains, it is possible that minor advances occurred and were overridden by subsequent advances during the LIA. Regardless, late glacial to Holocene moraines deposited in Alaska narrowly exceed LIA advances, indicating – without considering the role of precipitation – that late glacial and early to middle Holocene cooling events were minor in Alaska. While this observation may appear at odds with the magnitude of North Atlantic climate oscillations observed in Greenland Ice Core records (Fig. 7), mounting evidence suggests these oscillations may be more strongly expressed during winter compared to summer (e.g., Bromley et al., 2018). Because there is likely a heavy summer seasonal bias to alpine glacier-based temperature reconstructions, we suggest that predominantly wintertime North Atlantic oscillations would result in a somewhat dampened response from alpine glaciers, which is consistent with observed minor late glacial cooling at our site. Indeed, there is evidence from the southwestern Greenland Ice Sheet margin suggesting that the margin was retreating during the YD in response to summertime and local feedbacks, seemingly in contrast to the notable cooling observed in the Greenland ice core records (Funder et al., 2021; Carlson et

al., 2021). This observation might further support the notion of YD cooling being a predominantly winter phenomenon.

In the context of detailed glacier reconstructions observed elsewhere around the world, we find both similarities and differences between our record and others (Fig. 7). Broadly, Southern Hemisphere glacier records indicate a coupling with global CO₂ reflected in Antarctic ice core records, while glaciers in Alaska follow patterns of alpine glacier recession in the Alps and climate fluctuations recorded in Greenland ice cores (Fig. 7). Even if rising global CO₂ forced major recession following deposition of the 17 ka moraine (F2-IIIa) and any evidence was overridden by subsequent moraine deposition events at our site, that scenario would still require a large-magnitude re-advance during HS-1 in Alaska based on our record. While evidence for YD advances in Alaska extends beyond our site (Young et al., 2019), we acknowledge that within the resolution of ¹⁰Be dating, the late glacial moraine at our site may have been deposited at the time of the Antarctic cold reversal “break point” like records in the Southern Hemisphere and a site in Norway (Putnam et al., 2023). Despite uncertainty in the age assignment of one moraine at our site, we still observe a dampened climate response in Alaska at intervals generally tracking North Atlantic climate oscillations – cooling and moraine deposition at the culmination of HS-1, retreat through the Bølling warm period, and slight cooling during the early YD (Fig. 7) – with substantial deviation from records in the Southern Hemisphere, which all reinforces the notion that the North Atlantic signature of climate evolution through deglaciation extended out to Alaska.

6 Conclusions

New ¹⁰Be ages from late Holocene moraines in the Revelation Mountains reveal similar culminations with other, well-dated records in southern Alaska at ~1200–1300 CE and ~1700 CE, suggesting similar climatic controls across southern and western Alaska. The ELA-based temperature record from the Revelation Mountains, pinned by the late Holocene moraine constraints, indicates an LGM temperature depression of ~4 °C relative to the LIA, consistent with other limited proxy data from across the state. Most warming was delayed following global CO₂ rise until ~16 ka when abrupt ~1.5 °C of warming forced significant retreat. Brief intervals of cooling and moraine deposition late during HS-1 and in the early YD interrupted the overall pattern of warming and glacier retreat, similar to the observed glacial records in the Alps. Paired with ~2 °C of warming and glacier recession during the Bølling, these observations highlight the potential influence of North Atlantic forcing on climate in Alaska.

Data availability. All observations necessary to recalculate ¹⁰Be concentrations and exposure ages are reported in the main text, as well as in the ICE-D online database (<https://www.ice-d.org>, Balco, 2020).

Author contributions. JPT was responsible for conceptualization, formal analyses, visualizations, and writing of the original draft for this paper. JPB, NEY, and JMS provided resources, funding acquisition, conceptualization, and review and editing of the original draft.

Competing interests. The contact author has declared that none of the authors has any competing interests.

Disclaimer. Publisher’s note: Copernicus Publications remains neutral with regard to jurisdictional claims made in the text, published maps, institutional affiliations, or any other geographical representation in this paper. While Copernicus Publications makes every effort to include appropriate place names, the final responsibility lies with the authors.

Acknowledgements. We would like to acknowledge the fact that samples in this study were collected on ancestral lands of the Dënëneh and Dena’ina Elnena and that the University at Buffalo exists on the land of the Seneca. These peoples are the traditional caretakers of these lands. We give thanks for the opportunity to exist and work on lands that are rightfully theirs. We thank Roseanne Schwartz for lab processing and LLNL for ¹⁰Be measurements. Nicolás E. Young and Joerg M. Schaefer acknowledge support from the LDEO Climate Center.

Financial support. This research has been supported by the National Science Foundation (NSF award number 1853705).

Review statement. This paper was edited by Alberto Reyes and reviewed by two anonymous referees.

References

- Badding, M. E., Briner, J. P., and Kaufman, D. S.: ¹⁰Be ages of late Pleistocene deglaciation and Neoglaciation in the north-central Brooks Range, Arctic Alaska, *J. Quaternary Sci.*, 28, 95–102, <https://doi.org/10.1002/jqs.2596>, 2013.
- Balco, G.: Technical note: A prototype transparent-middle-layer data management and analysis infrastructure for cosmogenic-nuclide exposure dating, *Geochronology*, 2, 169–175, <https://doi.org/10.5194/gchron-2-169-2020>, 2020.
- Balco, G., Stone, J. O., Lifton, N. A., and Dunai, T. J.: A complete and easily accessible means of calculating surface exposure ages or erosion rates from ¹⁰Be and ²⁶Al measurements, *Quat. Geochronol.*, 3, 174–195, 2008.

- Balco, G., Briner, J., Finkel, R. C., Rayburn, J. A., Ridge, J. C., and Schaefer, J. M.: Regional beryllium-10 production rate calibration for late-glacial northeastern North America, *Quat. Geochronol.*, 4, 93–107, 2009.
- Barclay, D. J., Wiles, G. C., and Calkin, P. E.: Holocene glacier fluctuations in Alaska, *Quaternary Sci. Rev.*, 28, 2034–2048, <https://doi.org/10.1016/j.quascirev.2009.01.016>, 2009.
- Bereiter, B., Eggleston, S., Schmitt, J., Nehrbass-Ahles, C., Stocker, T. F., Fischer, H., Kipfstuhl, S., and Chappellaz, J.: Revision of the EPICA Dome C CO₂ record from 800 to 600 kyr before present, *Geophys. Res. Lett.*, 42, 542–549, 2015.
- Braithwaite, R. J.: Temperature and precipitation climate at the equilibrium-line altitude of glaciers expressed by the degree-day factor for melting snow, *J. Glaciol.*, 54, 437–444, <https://doi.org/10.3189/002214308785836968>, 2008.
- Briner, J. P., Kaufman, D. S., Manley, W. F., Finkel, R. C., and Caffee, M. W.: Cosmogenic exposure dating of late Pleistocene moraine stabilization in Alaska, *Geol. Soc. Am. Bull.*, 117, 1108–1120, 2005.
- Bromley, G., Putnam, A., Borns Jr., H., Lowell, T., Sandford, T., and Barrell, D.: Interstadial Rise and Younger Dryas Demise of Scotland's Last Ice Fields, *Paleoceanography and Paleoclimatology*, 33, 412–429, <https://doi.org/10.1002/2018PA003341>, 2018.
- Buizert, C., Gkinis, V., Severinghaus, J. P., He, F., Lecavalier, B. S., Kindler, P., Leuenberger, M., Carlson, A. E., Vinther, B., and Masson-Delmotte, V.: Greenland temperature response to climate forcing during the last deglaciation, *Science*, 345, 1177–1180, 2014.
- Carlson, A. E., Reyes, A. V., Gusterson, E., Axford, Y., Wilcken, K. M., and Rood, D. H.: Direct evidence for thinning and retreat of the southernmost Greenland ice sheet during the Younger Dryas, *Quaternary Sci. Rev.*, 267, 107105, <https://doi.org/10.1016/j.quascirev.2021.107105>, 2021.
- Claude, A., Ivy-Ochs, S., Kober, F., Antognini, M., Salcher, B., and Kubik, P. W.: The Chironico landslide (Valle Leventina, southern Swiss Alps): age and evolution, *Swiss J. Geosci.*, 107, 273–291, 2014.
- Daniels, W. C., Russell, J. M., Morrill, C., Longo, W. M., Giblin, A. E., Holland-Stergar, P., Welker, J. M., Wen, X., Hu, A., and Huang, Y.: Lacustrine leaf wax hydrogen isotopes indicate strong regional climate feedbacks in Beringia since the last ice age, *Quaternary Sci. Rev.*, 269, 107130, <https://doi.org/10.1016/j.quascirev.2021.107130>, 2021.
- Denton, G. H., Lowell, T., Heusser, C., Schlüchter, C., Andersen, B. G., Heusser, L. E., Moreno, P. I., and Marchant, D. R.: Geomorphology, stratigraphy, and radiocarbon chronology of Llanquihue Drift in the area of the Southern Lake District, Seno Reloncaví, and Isla Grande de Chiloé, Chile, *Geogr. Ann. A*, 81, 167–229, 1999.
- Dorfman, J., Stoner, J., Finkenbinder, M., Abbott, M., Xuan, C., and St-Onge, G.: A 37,000-year environmental magnetic record of aeolian dust deposition from Burial Lake, Arctic Alaska, *Quaternary Sci. Rev.*, 128, 81–97, 2015.
- Dortch, J. M., Owen, L. A., Caffee, M. W., and Brease, P.: Late Quaternary glaciation and equilibrium line altitude variations of the McKinley River region, central Alaska Range, Boreas, 39, 233–246, 2010.
- Ellis, J. M. and Calkin, P. E.: Chronology of Holocene glaciation, central Brooks range, Alaska, *Geol. Soc. Am. Bull.*, 95, 897–912, 1984.
- Finkenbinder, M., Abbott, M., Finney, B., Stoner, J., and Dorfman, J.: A multi-proxy reconstruction of environmental change spanning the last 37,000 years from Burial Lake, Arctic Alaska, *Quaternary Sci. Rev.*, 126, 227–241, 2015.
- Finkenbinder, M. S., Abbott, M. B., Edwards, M. E., Langdon, C. T., Steinman, B. A., and Finney, B. P.: A 31,000 year record of paleoenvironmental and lake-level change from Harding Lake, Alaska, USA, *Quaternary Sci. Rev.*, 87, 98–113, 2014.
- Funder, S., Sørensen, A. H. L., Larsen, N. K., Björk, A. A., Briner, J. P., Olsen, J., Schomacker, A., Levy, L. B., and Kjær, K. H.: Younger Dryas ice margin retreat in Greenland: new evidence from southwestern Greenland, *Clim. Past*, 17, 587–601, <https://doi.org/10.5194/cp-17-587-2021>, 2021.
- Heyman, J., Stroeven, A. P., Harbor, J. M., and Caffee, M. W.: Too young or too old: evaluating cosmogenic exposure dating based on an analysis of compiled boulder exposure ages, *Earth Planet. Sc. Lett.*, 302, 71–80, 2011.
- Hugonnet, R., McNabb, R., Berthier, E., Menounos, B., Nuth, C., Girod, L., Farinotti, D., Huss, M., Dussaillant, I., Brun, F., and Käab, A.: Accelerated global glacier mass loss in the early twenty-first century, *Nature*, 592, 726–731, <https://doi.org/10.1038/s41586-021-03436-z>, 2021.
- Hutchinson, M. F., Xu, T., and Stein, J. A.: Recent progress in the ANUDEM elevation gridding procedure, *Geomorphometry*, 2011, 19–22, 2011.
- Ivy-Ochs, S.: Glacier variations in the European Alps at the end of the last glaciation, *Cuadernos de investigación geográfica/Geographical Research Letters*, 295–315, 2015.
- Kaufman, D. S., Anderson, R. S., Hu, F. S., Berg, E., and Werner, A.: Evidence for a variable and wet Younger Dryas in southern Alaska, *Quaternary Sci. Rev.*, 29, 1445–1452, 2010.
- Kaufman, D. S., Young, N. E., Briner, J. P., and Manley, W. F.: Alaska Palaeo-Glacier Atlas (Version 2), in: *Developments in Quaternary Sciences*, vol. 15, edited by: Ehlers, J., Gibbard, P. L., and Hughes, P. D., Elsevier, 427–445, <https://doi.org/10.1016/B978-0-444-53447-7.00033-7>, 2011.
- Kern, Z. and László, P.: Size specific steady-state accumulation-area ratio: an improvement for equilibrium-line estimation of small palaeoglaciers, *Quaternary Sci. Rev.*, 29, 2781–2787, 2010.
- Kobashi, T., Menviel, L., Jeltsch-Thömmes, A., Vinther, B. M., Box, J. E., Muscheler, R., Nakaegawa, T., Pfister, P. L., Döring, M., and Leuenberger, M.: Volcanic influence on centennial to millennial Holocene Greenland temperature change, *Scientific reports*, 7, 1441, <https://doi.org/10.1038/s41598-017-01451-7>, 2017.
- Kurek, J., Cwynar, L. C., Ager, T. A., Abbott, M. B., and Edwards, M. E.: Late Quaternary paleoclimate of western Alaska inferred from fossil chironomids and its relation to vegetation histories, *Quaternary Sci. Rev.*, 28, 799–811, 2009.
- Lal, D.: Cosmic ray labeling of erosion surfaces: in situ nuclide production rates and erosion models, *Earth Planet. Sc. Lett.*, 104, 424–439, 1991.
- Matmon, A., Briner, J., Carver, G., Bierman, P., and Finkel, R.: Moraine chronosequence of the Donnelly Dome region, Alaska, *Quaternary Res.*, 74, 63–72, 2010.

- Nishiizumi, K., Imamura, M., Caffee, M. W., Southon, J. R., Finkel, R. C., and McAninch, J.: Absolute calibration of ^{10}Be AMS standards, *Nuclear Instruments and Methods in Physics Research Section B: Beam Interactions with Materials and Atoms*, 258, 403–413, <https://doi.org/10.1016/j.nimb.2007.01.297>, 2007.
- Oerlemans J.: Extracting a Climate Signal from 169 Glacier Records, *Science*, 308, 675–677, <https://doi.org/10.1126/science.1107046>, 2005.
- Ohmura, A., Kasser, P., and Funk, M.: Climate at the Equilibrium Line of Glaciers, *J. Glaciol.*, 38, 397–411, <https://doi.org/10.3189/S002214300002276>, 1992.
- Osmaston, H.: Estimates of glacier equilibrium line altitudes by the Area \times Altitude, the Area \times Altitude Balance Ratio and the Area \times Altitude Index methods and their validation, *Quatern. Int.*, 138–139, 22–31, <https://doi.org/10.1016/j.quaint.2005.02.004>, 2005.
- Palacios, D., Stokes, C. R., Phillips, F. M., Clague, J. J., Alcalá-Reygosa, J., Andrés, N., Angel, I., Blard, P.-H., Briner, J. P., and Hall, B. L.: The deglaciation of the Americas during the Last Glacial Termination, *Earth-Sci. Rev.*, 203, 103113, <https://doi.org/10.1016/j.earsci.2020.103113>, 2020.
- Pellitero, R., Rea, B. R., Spagnolo, M., Bakke, J., Hughes, P., Ivy-Ochs, S., Lukas, S., and Ribolini, A.: A GIS tool for automatic calculation of glacier equilibrium-line altitudes, *Comput. Geosci.*, 82, 55–62, <https://doi.org/10.1016/j.cageo.2015.05.005>, 2015.
- Pellitero, R., Rea, B. R., Spagnolo, M., Bakke, J., Ivy-Ochs, S., Frew, C. R., Hughes, P., Ribolini, A., Lukas, S., and Renssen, H.: GlaRe, a GIS tool to reconstruct the 3D surface of palaeoglaciers, *Comput. Geosci.*, 94, 77–85, <https://doi.org/10.1016/j.cageo.2016.06.008>, 2016.
- Pendleton, S. L., Briner, J. P., Kaufman, D. S., and Zimmerman, S. R.: Using Cosmogenic ^{10}Be Exposure Dating and Lichenometry to Constrain Holocene Glaciation in the Central Brooks Range, Alaska, *Arct. Antarct. Alp. Res.*, 49, 115–132, <https://doi.org/10.1657/AAAR0016-045>, 2017.
- Praetorius, S. K., Condrion Alan, Mix Alan C., Walczak Maureen H., McKay Jennifer L., and Du Jianghui: The role of Northeast Pacific meltwater events in deglacial climate change, *Sci. Adv.*, 6, eaay2915, <https://doi.org/10.1126/sciadv.aay2915>, 2020.
- Putnam, A. E., Schaefer, J. M., Denton, G. H., Barrell, D. J., Birkel, S. D., Andersen, B. G., Kaplan, M. R., Finkel, R. C., Schwartz, R., and Doughty, A. M.: The last glacial maximum at 44° S documented by a ^{10}Be moraine chronology at Lake Ohau, Southern Alps of New Zealand, *Quaternary Sci. Rev.*, 62, 114–141, 2013.
- Putnam, A. E., Bromley, G. R., Rademaker, K., and Schaefer, J. M.: In situ ^{10}Be production-rate calibration from a ^{14}C -dated late-glacial moraine belt in Rannoch Moor, central Scottish Highlands, *Quat. Geochronol.*, 50, 109–125, 2019.
- Putnam, A. E., Denton, G. H., and Schaefer, J. M.: A ^{10}Be chronology of the Esmark Moraine and Lysefjorden region, southwestern Norway: Evidence for coeval glacier resurgence in both polar hemispheres during the Antarctic Cold Reversal, *Quaternary Sci. Rev.*, 316, 108259, <https://doi.org/10.1016/j.quascirev.2023.108259>, 2023.
- Roe, G. H., Baker, M. B., and Herla, F.: Centennial glacier retreat as categorical evidence of regional climate change, *Nat. Geosci.*, 10, 95–99, <https://doi.org/10.1038/ngeo2863>, 2017.
- Rupper, S. and Roe, G.: Glacier changes and regional climate: a mass and energy balance approach, *J. Climate*, 21, 5384–5401, 2008.
- Schaefer, J. M., Denton, G. H., Barrell, D. J., Ivy-Ochs, S., Kubik, P. W., Andersen, B. G., Phillips, F. M., Lowell, T. V., and Schlüchter, C.: Near-synchronous interhemispheric termination of the last glacial maximum in mid-latitudes, *Science*, 312, 1510–1513, 2006.
- Schaefer, J. M., Denton, G. H., Kaplan, M., Putnam, A., Finkel, R. C., Barrell, D. J., Andersen, B. G., Schwartz, R., Mackintosh, A., and Chinn, T.: High-frequency Holocene glacier fluctuations in New Zealand differ from the northern signature, *Science*, 324, 622–625, 2009.
- Stone, J. O.: Air pressure and cosmogenic isotope production, *J. Geophys. Res.-Sol. Ea.*, 105, 23753–23759, 2000.
- Strelin, J., Denton, G., Vandergoes, M., Ninnemann, U., and Putnam, A.: Radiocarbon chronology of the late-glacial Puerto Bandera moraines, southern Patagonian Icefield, Argentina, *Quaternary Sci. Rev.*, 30, 2551–2569, 2011.
- Tierney, J. E., Zhu, J., King, J., Malevich, S. B., Hakim, G. J., and Poulsen, C. J.: Glacial cooling and climate sensitivity revisited, *Nature*, 584, 569–573, 2020.
- Tulenko, J. P., Briner, J. P., Young, N. E., and Schaefer, J. M.: Beryllium-10 chronology of early and late Wisconsinan moraines in the Revelation Mountains, Alaska: Insights into the forcing of Wisconsinan glaciation in Beringia, *Quaternary Sci. Rev.*, 197, 129–141, <https://doi.org/10.1016/j.quascirev.2018.08.009>, 2018.
- Tulenko, J. P., Briner, J. P., Young, N. E., and Schaefer, J. M.: The last deglaciation of Alaska and a new benchmark ^{10}Be moraine chronology from the western Alaska Range, *Quaternary Sci. Rev.*, 287, 107549, 2022.
- Valentino, J. D., Owen, L. A., Spotila, J. A., Cesta, J. M., and Caffee, M. W.: Timing and extent of Late Pleistocene glaciation in the Chugach Mountains, Alaska, *Quaternary Res.*, 101, 205–224, 2021.
- Verbyla, D. and Kurkowski, T. A.: NDVI–Climate relationships in high-latitude mountains of Alaska and Yukon Territory, *Arct. Antarct. Alp. Res.*, 51, 397–411, <https://doi.org/10.1080/15230430.2019.1650542>, 2019.
- Viau, A., Gajewski, K., Sawada, M., and Bunbury, J.: Low- and high-frequency climate variability in eastern Beringia during the past 25 000 years, *Can. J. Earth Sci.*, 45, 1435–1453, 2008.
- Walcott, C. K., Briner, J. P., Tulenko, J. P., and Evans, S. M.: Equilibrium line altitudes of alpine glaciers in Alaska suggest Last Glacial Maximum summer temperature was 2–5 °C lower than during the pre-industrial, *Clim. Past*, 20, 91–106, <https://doi.org/10.5194/cp-20-91-2024>, 2024.
- Wiles, G. C., Jacoby, G. C., Davi, N. K., and McAllister, R. P.: Late Holocene glacier fluctuations in the Wrangell Mountains, Alaska, *Geol. Soc. Am. Bull.*, 114, 896–908, 2002.
- Wiles, G. C., D’Arrigo, R. D., Villalba, R., Calkin, P. E., and Barclay, D. J.: Century-scale solar variability and Alaskan temperature change over the past millennium, *Geophys. Res. Lett.*, 31, L15203, <https://doi.org/10.1029/2004GL020050>, 2004.
- Wittmeier, H. E., Schaefer, J. M., Bakke, J., Rupper, S., Paasche, Ø., Schwartz, R., and Finkel, R. C.: Late Glacial mountain glacier culmination in Arctic Norway prior to

- the Younger Dryas, *Quaternary Sci. Rev.*, 245, 106461, <https://doi.org/10.1016/j.quascirev.2020.106461>, 2020.
- Young, N. E., Schaefer, J. M., Briner, J. P., and Goehring, B. M.: A ^{10}Be production-rate calibration for the Arctic, *J. Quaternary Sci.*, 28, 515–526, 2013.
- Young, N. E., Briner, J. P., Schaefer, J., Zimmerman, S., and Finkel, R. C.: Early Younger Dryas glacier culmination in southern Alaska: Implications for North Atlantic climate change during the last deglaciation, *Geology*, 47, 550–554, 2019.